

## WEST Search History

DATE: Tuesday, February 15, 2005

Hide?	Set Name	Query	Hit Count
	<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=AND</i>		
<input type="checkbox"/>	L1	clostrid\$.clm. and perfring\$.clm.	168
<input type="checkbox"/>	L2	L1 and promoter.clm.	9
<input type="checkbox"/>	L3	L1 and promoter.clm.	9
<input type="checkbox"/>	L4	clostrid\$ near6 perfring\$	2854
<input type="checkbox"/>	L5	L4 near50 promoter	4
<input type="checkbox"/>	L6	L5 not l3	3

END OF SEARCH HISTORY

- 
- ☐ 1. 20040029201. 11 Jul 03. 12 Feb 04. Identification of targets of antimicrobial compounds. Fan, Frank, et al. 435/7.23; G01N033/574.
- 
- ☐ 2. 20040029129. 25 Oct 02. 12 Feb 04. Identification of essential genes in microorganisms. Wang, Liangsu, et al. 435/6; 435/183 435/252.33 435/254.2 435/320.1 435/325 435/419 435/69.1 530/350 536/23.2 C12Q001/68 C07H021/04 C12N001/20 C12N009/00 C12P021/02 C12N001/21 C07K014/47 C12N005/04 C12N001/18.
- 
- ☐ 3. 20030186281. 20 Dec 02. 02 Oct 03. Modified tetracycline repressor protein compositions and methods of use. Hillen, Wolfgang. 435/6; 435/191 435/252.3 435/252.31 435/320.1 435/69.1 435/7.32 536/23.2 C12Q001/68 G01N033/554 G01N033/569 C07H021/04 C12N009/06 C12N001/21 C12P021/02 C12N015/74.
- 
- ☐ 4. 20030124507. 13 Mar 02. 03 Jul 03. Method of generating conditionally expressed mutant cells using expressible antisense sequences. Marra, Andrea, et al. 435/4; 435/252.3 435/471 435/6 C12Q001/00 C12Q001/68 C12N015/74 C12N001/21.
- 
- ☐ 5. 20030027286. 21 Dec 01. 06 Feb 03. Bacterial promoters and methods of use. Haselbeck, Robert, et al. 435/69.6; 435/219 435/252.3 435/320.1 435/6 536/23.2 C07H021/04 C12P021/04 C12N009/50 C12Q001/68 C12N001/21 C12N015/74.
- 
- ☐ 6. 20020086310. 30 Aug 01. 04 Jul 02. Identification of targets of antimicrobial compounds. Fan, Frank, et al. 435/6; 435/32 C12Q001/68 C12Q001/18.
- 
- ☐ 7. 6780447. 19 Feb 02; 24 Aug 04. Bacteriocin-containing sorbic acid product as addition to feedstuffs in agricultural livestock rearing. Raczek, Nico N.. 426/61; A23K001/18.
- 
- ☐ 8. 5955368. 06 Apr 98; 21 Sep 99. Expression system for clostridium species. Johnson, Eric A., et al. 435/488; 435/252.3 435/320.1 435/476 536/23.1 536/24.1. C12N001/21 C12N015/70 C12N015/74 C12N015/64.
- 
- ☐ 9. 5004692. 15 Dec 87; 02 Apr 91. Cloning and expression of phospholipase C genes. Tso, J. Yun, et al. 435/183; 435/195 435/252.3 435/252.33 435/320.1 435/358 435/365 435/367 536/23.2 536/23.7. C12N009/00 C12N009/14.

- 
- ☐ 1. 20040029201. 11 Jul 03. 12 Feb 04. Identification of targets of antimicrobial compounds. Fan, Frank, et al. 435/7.23; G01N033/574.
- 
- ☐ 2. 20040029129. 25 Oct 02. 12 Feb 04. Identification of essential genes in microorganisms. Wang, Liangsu, et al. 435/6; 435/183 435/252.33 435/254.2 435/320.1 435/325 435/419 435/69.1 530/350 536/23.2 C12Q001/68 C07H021/04 C12N001/20 C12N009/00 C12P021/02 C12N001/21 C07K014/47 C12N005/04 C12N001/18.
- 
- ☐ 3. 20030186281. 20 Dec 02. 02 Oct 03. Modified tetracycline repressor protein compositions and methods of use. Hillen, Wolfgang. 435/6; 435/191 435/252.3 435/252.31 435/320.1 435/69.1 435/7.32 536/23.2 C12Q001/68 G01N033/554 G01N033/569 C07H021/04 C12N009/06 C12N001/21 C12P021/02 C12N015/74.
- 
- ☐ 4. 20030124507. 13 Mar 02. 03 Jul 03. Method of generating conditionally expressed mutant cells using expressible antisense sequences. Marra, Andrea, et al. 435/4; 435/252.3 435/471 435/6 C12Q001/00 C12Q001/68 C12N015/74 C12N001/21.
- 
- ☐ 5. 20030027286. 21 Dec 01. 06 Feb 03. Bacterial promoters and methods of use. Haselbeck, Robert, et al. 435/69.6; 435/219 435/252.3 435/320.1 435/6 536/23.2 C07H021/04 C12P021/04 C12N009/50 C12Q001/68 C12N001/21 C12N015/74.
- 
- ☐ 6. 20020086310. 30 Aug 01. 04 Jul 02. Identification of targets of antimicrobial compounds. Fan, Frank, et al. 435/6; 435/32 C12Q001/68 C12Q001/18.
- 
- ☐ 7. 6780447. 19 Feb 02; 24 Aug 04. Bacteriocin-containing sorbic acid product as addition to feedstuffs in agricultural livestock rearing. Raczek, Nico N.. 426/61;. A23K001/18.
- 
- ☐ 8. 5955368. 06 Apr 98; 21 Sep 99. Expression system for clostridium species. Johnson; Eric A., et al. 435/488; 435/252.3 435/320.1 435/476 536/23.1 536/24.1. C12N001/21 C12N015/70 C12N015/74 C12N015/64.
- 
- ☐ 9. 5004692. 15 Dec 87; 02 Apr 91. Cloning and expression of phospholipase C genes. Tso; J. Yun, et al. 435/183; 435/195 435/252.3 435/252.33 435/320.1 435/358 435/365 435/367 536/23.2 536/23.7. C12N009/00 C12N009/14.
-

[First Hit](#)      [Previous Doc](#)      [Next Doc](#)      [Go to Doc#](#)

End of Result Set



Generate Collection

Print

L6: Entry 3 of 3

File: DWPI

Mar 6, 2003

DERWENT-ACC-NO: 1999-217498

DERWENT-WEEK: 200433

COPYRIGHT 2005 DERWENT INFORMATION LTD

TITLE: Clostridium beta2 toxin gene promoter and signal sequence - useful against toxins from Clostridium perfringens

Basic Abstract Text (1):

The nucleic acid of the Clostridium perfringens beta 2 toxin gene promoter comprising at least part of sequence (I) (given in the specification), is new.

ATTTGGGATA TCTTAAATTT AGCACAGAAG AATGTTTAAA TGAAATAAAG ATAATAAAAA GATATATTAA  
TTATATAGCT GAAAATTTAT AATTATATGA TAAGTATAGT TAATAAATAA AAAGTGTCT CGGGGGACAC  
TTTTTTGTTT TAAAAAGGAA AATATAAATA AAATTTAGAT AAAAGTGTA AATAATTATT TTTATTTTAA  
ATTTGTTAAA AATTTGATAT AATTGAATTG TAAAAAAAAT TTCAGGGGGG AATATAAATG AAAATTATTA  
TTTCAAAGTT TACTGTAATT TTTATGTTTT CATGTTTCT TATTGTT (I). NOTE: The last 60 bases of  
(I) encode the first 20 amino acids of the signal peptide (II).

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

File 53:FOODLINE(R): Science Sight 1972-2005/Feb 14  
 (c) 2005 LFRA  
 \*File 53: Price change effective Jan 1, 2005. Enter HELP  
 RATES 53 for details.  
 File 149:TGG Health&Wellness DB(SM) 1976-2005/Feb W1  
 (c) 2005 The Gale Group  
 File 203:AGRIS 1974-2004/Nov  
 Dist by NAL, Intl Copr. All rights reserved  
 File 292:GEOBASE(TM) 1980-2005/Jan B1  
 (c) 2005 Elsevier Science Ltd.  
 \*File 292: Price change effective Jan 1, 2005. Enter HELP  
 RATES 292 for details.  
 File 342:Derwent Patents Citation Indx 1978-05/200506  
 (c) 2005 Thomson Derwent  
 File 345:Inpadoc/Fam.& Legal Stat 1968-2004/UD=200504  
 (c) 2005 EPO

Set Items Description

Cost is in DialUnits  
 ?ds

Set	Items	Description
S1	203	(CLOSTRID? OR PERFRINGEN?)/TI AND PROMOTER?/TI
S2	75	RD (unique items)
?t s2/9/6	12 13 14 15 16 17 23 66 67 68	

2/9/6 (Item 6 from file: 155)  
 DIALOG(R) File 155:MEDLINE(R)  
 (c) format only 2005 The Dialog Corp. All rts. reserv.

13159543 PMID: 8828224

**An upstream activating sequence containing curved DNA involved in activation of the Clostridium perfringens plc promoter .**

Matsushita C; Matsushita O; Katayama S; Minami J; Takai K; Okabe A  
 Department of Microbiology, Kagawa Medical School, Japan.  
 Microbiology (Reading, England) (ENGLAND) Sep 1996, 142 ( Pt 9)  
 p2561-6, ISSN 1350-0872 Journal Code: 9430468  
 Document type: Journal Article  
 Languages: ENGLISH  
 Main Citation Owner: NLM  
 Record type: Completed  
 Subfile: INDEX MEDICUS

The plc gene, which encodes phospholipase C (alpha-toxin) of Clostridium perfringens, possesses three poly(A) tracts forming an intrinsically curved DNA region immediately upstream of the promoter. The in vivo transcriptional activity of the plasmid-borne plc gene was stimulated by this curved-DNA-containing sequence, depending on its proper linear and rotational orientation. The in vitro transcriptional activity of the plc gene was also stimulated by the upstream sequence. In addition, the stimulatory effect of the sequence and the degree of DNA bending were greater at lower temperature, as was demonstrated by both in vitro and in vivo transcription assays, and a gel-mobility assay, respectively. A similar temperature effect was also observed with the chromosomal plc gene. These observations suggest that the upstream DNA curvature per se stimulates the initiation of transcription of the plc gene, possibly through direct contact with RNA polymerase.

Tags: Support, Non-U.S. Gov't

Descriptors: \*Clostridium perfringens--genetics--GE; \*Phospholipase C  
 --genetics--GE; Base Sequence; Chromosome Mapping; Chromosomes--genetics  
 --GE; Chromosomes--physiology--PH; DNA--physiology--PH; Gene Expression  
 Regulation, Bacterial; Molecular Sequence Data; Mutagenesis, Insertional;  
 Mutagenesis, Site-Directed; Nucleic Acid Conformation; Plasmids--genetics  
 --GE; Plasmids--physiology--PH; Promoter Regions (Genetics); Sequence  
 Deletion; Temperature; Transcription, Genetic

CAS Registry No.: 0 (Plasmids); 9007-49-2 (DNA)

Enzyme No.: EC 3.1.4.3 (Phospholipase C)

Record Date Created: 19970113  
Record Date Completed: 19970113

2/9/12 (Item 12 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2005 The Dialog Corp. All rts. reserv.

10260070 PMID: 7960138

**Expression from the Clostridium perfringens cpe promoter in C. perfringens and Bacillus subtilis.**

Melville S B; Labbe R; Sonenshein A L

Department of Molecular Biology and Microbiology, Tufts University School of Medicine, Boston, Massachusetts 02111.

Infection and immunity (UNITED STATES) Dec 1994, 62 (12) p5550-8,  
ISSN 0019-9567 Journal Code: 0246127

Contract/Grant No.: GM42219; GM; NIGMS; P30 DK34928; DK; NIDDK

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

Clostridium perfringens is a source of food poisoning in humans and animals because of production of a potent enterotoxin (CPE). To study the regulation of the cpe gene in C. perfringens, we cloned and sequenced the cpe promoter regions and N-terminal domains from three strains. The cpe promoter region from one strain contained a 45-bp insertion compared with previously published sequences. This insertion was also found in two (of five) other Cpe+ strains. cpe gene expression in C. perfringens was measured by using translational fusions of each promoter type to the Escherichia coli gusA gene, which codes for beta-glucuronidase. For either promoter type, cpe-gusA expression was undetectable throughout exponential growth but increased dramatically at the beginning of the stationary phase. To measure cpe expression in Bacillus subtilis, cpe-gusA fusions were integrated into the B. subtilis chromosome. Both types of promoter exhibited moderate expression during exponential growth; cpe expression increased threefold at the beginning of the stationary phase. Transcriptional start sites were determined by primer extension and in vitro transcription assays. For C. perfringens, both types of promoter gave the same 5' end, 197 bp upstream of the translation start (50 bp downstream of the 45-bp insertion). In B. subtilis, however, the 5' end was internal to the 45-bp insertion, suggesting the use of a different promoter than that utilized by C. perfringens.

Tags: Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: \*Clostridium perfringens--genetics--GE; \*Enterotoxins--genetics--GE; \*Gene Expression Regulation, Bacterial; \*Promoter Regions (Genetics)--genetics--GE; Bacillus subtilis--genetics--GE; Base Sequence; Cell-Free System; Cloning, Molecular; Clostridium perfringens--growth and development--GD; DNA, Recombinant; Electroporation; Enterotoxins--biosynthesis--BI; Glucuronidase--biosynthesis--BI; Glucuronidase--genetics--GE; Molecular Sequence Data; Polymerase Chain Reaction; RNA, Messenger--genetics--GE; Raffinose--pharmacology--PD; Recombinant Fusion Proteins--biosynthesis--BI; Sequence Analysis, DNA; Spores, Bacterial--drug effects--DE; Spores, Bacterial--growth and development--GD; Starch--pharmacology--PD; Transcription, Genetic; Translation, Genetic  
Molecular Sequence Databank No.: GENBANK/U11257; GENBANK/U11259; GENBANK/U11294

CAS Registry No.: 0 (DNA, Recombinant); 0 (Enterotoxins); 0 (RNA, Messenger); 0 (Recombinant Fusion Proteins); 0 (enterotoxin, Clostridium); 512-69-6 (Raffinose); 9005-25-8 (Starch)

Enzyme No.: EC 3.2.1.31 (Glucuronidase)

Gene Symbol: cpe

Record Date Created: 19941229

Record Date Completed: 19941229

2/9/13 (Item 13 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2005 The Dialog Corp. All rts. reserv.

10170067 PMID: 8058826

**A Clostridium perfringens vector for the selection of promoters .**

Matsushita C; Matsushita O; Koyama M; Okabe A

Department of Microbiology, Kagawa Medical School, Japan.

Plasmid (UNITED STATES) May 1994, 31 (3) p317-9, ISSN 0147-619X

Journal Code: 7802221

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

A promoter selection vector for Clostridium perfringens genes was constructed from a C. perfringens-Escherichia coli shuttle vector, pJIR418. The plasmid carries a promoterless chloramphenicol acetyltransferase gene (catP), derived from pIP401, downstream of the multiple cloning sites of pUC18. When a promoter region of the phospholipase C gene was inserted into one of the cloning sites, derivatives of C. perfringens strain 13 carrying the resultant plasmid acquired resistance to chloramphenicol. This plasmid should be useful reporter system for C. perfringens genes.

Tags: Support, Non-U.S. Gov't

Descriptors: \*Clostridium perfringens--genetics--GE; \*Genetic Vectors; \*Plasmids; \*Promoter Regions (Genetics); Base Sequence; Blotting, Northern; Chloramphenicol O-Acetyltransferase--biosynthesis--BI; Chloramphenicol O-Acetyltransferase--genetics--GE; Chloramphenicol O-Acetyltransferase--metabolism--ME; Cloning, Molecular--methods--MT; DNA Primers; Escherichia coli; Genes, Bacterial; Molecular Sequence Data; RNA, Messenger--analysis--AN; RNA, Messenger--biosynthesis--BI; Restriction Mapping

CAS Registry No.: 0 (DNA Primers); 0 (Genetic Vectors); 0 (Plasmids); 0 (RNA, Messenger)

Enzyme No.: EC 2.3.1.28 (Chloramphenicol O-Acetyltransferase)

Gene Symbol: catP

Record Date Created: 19940914

Record Date Completed: 19940914

2/9/14 (Item 14 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2005 The Dialog Corp. All rts. reserv.

08801833 PMID: 1981087

**Studies on Clostridium acetobutylicum glnA promoters and antisense RNA.**

Janssen P J; Jones D T; Woods D R

Department of Microbiology, University of Cape Town, Rondebosch, South Africa.

Molecular microbiology (ENGLAND) Sep 1990, 4 (9) p1575-83, ISSN 0950-382X Journal Code: 8712028

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

The Clostridium acetobutylicum glnA gene has two transcript start sites under the control of promoters p1 and p2. Initiation of transcription was regulated by nitrogen and a downstream region was implicated in the regulation of transcript initiation by nitrogen in Escherichia coli. Putative antisense RNA was produced from a single downstream transcript start site under the control of p3. An up-promoter mutation in p3 resulted in lower levels of glutamine synthetase (GS) activity. Putative antisense RNA had a role in down-regulating GS expression but was not involved in regulation by nitrogen. Deletion of downstream inverted repeat sequences resulted in very low levels of GS activity.

Descriptors: \*Clostridium--genetics--GE; \*Gene Expression Regulation, Bacterial; \*Glutamate-Ammonia Ligase--genetics--GE; \*Promoter Regions

(Genetics); \*RNA, Antisense--genetics--GE; Base Sequence; Clostridium  
--enzymology--EN; Escherichia coli--genetics--GE; Genes, Bacterial;  
Glutamate-Ammonia Ligase--metabolism--ME; Molecular Sequence Data; Mutation  
; Nitrogen--pharmacology--PD; RNA, Antisense--metabolism--ME; Repetitive  
Sequences, Nucleic Acid; Transcription, Genetic

CAS Registry No.: 0 (RNA, Antisense); 7727-37-9 (Nitrogen)

Enzyme No.: EC 6.3.1.2 (Glutamate-Ammonia Ligase)

Gene Symbol: glnA

Record Date Created: 19910325

Record Date Completed: 19910325

2/9/15 (Item 15 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2005 The Dialog Corp. All rts. reserv.

07999446 PMID: 2463955

**Identification and characterization of Clostridium difficile promoter element that is functional in Escherichia coli.**

Dailey D C; Schloemer R H

Department of Microbiology and Immunology, Indiana University School of Medicine, Indianapolis 46223.

Gene (NETHERLANDS) Oct 30 1988, 70 (2) p343-50, ISSN 0378-1119

Journal Code: 7706761

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

The promoter element involved in the expression of a previously characterized cloned clostridial antigen was isolated and characterized. A restriction fragment containing the promoter element of the Clostridium difficile insert was cloned using the promoter probe vector, pGA46. Subclones of the clostridial DNA insert in pGA46 were then analyzed by nucleotide sequencing and by S1 nuclease experiments. The clostridial promoter element exhibits a high degree of homology with typical Escherichia coli promoter elements. This sequence probably represents a unique class of clostridial promoter elements which, given their ability to function in E. coli and C. difficile, can be used in the construction of a shuttle vector capable of gene expression in E. coli and C. difficile.

Descriptors: \*Clostridium--genetics--GE; \*Escherichia coli--genetics--GE; \*Promoter Regions (Genetics); \*Transformation, Genetic; Aspergillus Nuclease S1; Base Sequence; Blotting, Southern; Cloning, Molecular; Electrophoresis, Agar Gel; Endonucleases--diagnostic use--DU; Gene Expression Regulation; Genetic Vectors; Plasmids; RNA, Bacterial--isolation and purification--IP; Restriction Mapping; Tetracycline Resistance --genetics--GE

Molecular Sequence Databank No.: GENBANK/M22864

CAS Registry No.: 0 (Genetic Vectors); 0 (Plasmids); 0 (RNA, Bacterial)

Enzyme No.: EC 3.1.- (Endonucleases); EC 3.1.30.1 (Aspergillus Nuclease S1)

Record Date Created: 19890302

Record Date Completed: 19890302

2/9/16 (Item 16 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2005 The Dialog Corp. All rts. reserv.

07931034 PMID: 2460717

**Studies of UV-inducible promoters from Clostridium perfringens in vivo and in vitro.**

Garnier T; Cole S T

Biochimie des Regulations Cellulaires, Institut Pasteur, Paris, France.

Molecular microbiology (ENGLAND) Sep 1988, 2 (5) p607-14, ISSN

0950-382X Journal Code: 8712028



Document type: Journal Article  
Languages: ENGLISH  
Main Citation Owner: NLM  
Record type: Completed  
Subfile: INDEX MEDICUS

Expression of a 4 kb segment of the bacteriocinogenic plasmid, pIP404, from *Clostridium perfringens* is inducible by UV-irradiation. DNA sequence analysis revealed that this region contains three genes: *uviA*, *uviB* and *bcn* encoding the bacteriocin BCN5. Biochemical studies with mRNAs showed that expression was controlled at the transcriptional level and that the genes were organized in two independent transcriptional units, *uviAB* and *bcn*, both directed by tandem promoters inducible by UV light. The *bcn* gene is transcribed from three promoters (P1, P2, P3) while transcription of *uviAB* is directed by two promoters (P4, P5). With the exception of P4, which bears some resemblance to the consensus eubacterial promoter sequence, none of these promoters was recognized in vitro by the major forms of RNA polymerase from *C. perfringens*, *Bacillus subtilis* or *Escherichia coli*. Promoters P1, P3 and P5, which show striking homology with each other, contain unusual sequences in the '-35' and '-10' regions known to be recognized by RNA polymerase and this might indicate positive control.

Tags: Support, Non-U.S. Gov't

Descriptors: \**Clostridium perfringens*--genetics--GE; \*Gene Expression Regulation; \*Promoter Regions (Genetics); Amino Acid Sequence; Base Sequence; *Clostridium perfringens*--enzymology--EN; DNA-Directed RNA Polymerases--biosynthesis--BI; DNA-Directed RNA Polymerases--isolation and purification--IP; Molecular Sequence Data; Plasmids; RNA, Bacterial--biosynthesis--BI; RNA, Messenger--biosynthesis--BI; Restriction Mapping; Ultraviolet Rays

Molecular Sequence Databank No.: GENBANK/J03309; GENBANK/J03310

CAS Registry No.: 0 (Plasmids); 0 (RNA, Bacterial); 0 (RNA, Messenger)

Enzyme No.: EC 2.7.7.6 (DNA-Directed RNA Polymerases)

Record Date Created: 19881220

Record Date Completed: 19881220

2/9/17 (Item 17 from file: 155)

DIALOG(R)File 155:MEDLINE(R)

(c) format only 2005 The Dialog Corp. All rts. reserv.

07187369 PMID: 3733758

**In vivo and in vitro transcription of the *Clostridium pasteurianum* ferredoxin gene. Evidence for "extended" promoter elements in gram-positive organisms.**

Graves M C; Rabinowitz J C

Journal of biological chemistry (UNITED STATES) Aug 25 1986, 261 (24) p11409-15, ISSN 0021-9258 Journal Code: 2985121R

Contract/Grant No.: AI6712; AI; NIAID; AM2109-28; AM; NIADDK

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

Analysis of *Clostridium pasteurianum* genomic DNA indicates that the ferredoxin (Fd) gene is present in a single copy. The cloned Fd gene previously described (Graves, M.C., Mullenbach, G. T., and Rabinowitz, J. C. (1985) Proc. Natl. Acad. Sci. U. S. A. 82, 1653-1657) was used to map in vivo and in vitro synthesized Fd transcripts. The in vivo mRNA was sized in two ways: by Northern hybridization analysis, and more directly from the known DNA sequence after the 5'- and 3'-termini were identified. The 5'-end was determined by primer extension-dideoxy sequencing and the 3'-end by S1 nuclease mapping. The monocistronic Fd mRNA contains about 255 nucleotides and, thus, is one of the shortest bacterial mRNAs yet described. We also examined the Fd transcripts produced by *Escherichia coli* transformed with the plasmid containing the Fd gene. *E. coli* RNA polymerase most likely recognizes the same promoter (P1) as the clostridial polymerase, and furthermore, efficiently uses an additional promoter (P2) that is poorly

recognized by the normal host enzyme. For comparison, in vitro transcripts were generated by E. coli and Bacillus subtilis RNA polymerases. In vitro, only promoter P1 is used by either E. coli or B. subtilis RNA polymerase. The 3'-end of each of the four types of transcripts occurs essentially at the same location and maps to within a large dyad symmetry element. Comparison of the Fd promoter with other Gram-positive promoters reveals that some sequences outside of the traditional Pribnow and -35 regions are conserved. This analysis indicates that an "extended" promoter recognition site may be required in these organisms.

Tags: Support, U.S. Gov't, P.H.S.

Descriptors: \*Clostridium--genetics--GE; \*Ferredoxins--genetics--GE; \*Promoter Regions (Genetics); \*Transcription, Genetic; Base Sequence; Electrophoresis, Polyacrylamide Gel; Nucleic Acid Conformation; Nucleic Acid Hybridization; RNA, Messenger--metabolism--ME

Molecular Sequence Databank No.: GENBANK/M11214; GENBANK/M13633; GENBANK/M13682

CAS Registry No.: 0 (Ferredoxins); 0 (RNA, Messenger)

Record Date Created: 19860919

Record Date Completed: 19860919

2/9/23 (Item 6 from file: 5)

DIALOG(R) File 5: Biosis Previews(R)

(c) 2005 BIOSIS. All rts. reserv.

0010820891 BIOSIS NO.: 199799454951

**The level of expression of alpha-toxin by different strains of Clostridium perfringens is dependent on differences in promoter structure and genetic background**

AUTHOR: Bullifent Helen L; Moir Anne; Awad Milna M; Scott Paul T; Rood Julian I; Titball Richard W

AUTHOR ADDRESS: Defence Evaluation and Res. Agency, CBD Porton Down, Salisbury, Wiltshire SP4 0JQ, UK\*\*UK

JOURNAL: Anaerobe 2 (6): p365-371 1996 1996

ISSN: 1075-9964

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The control of expression of the alpha-toxin gene (cpa or plc) of Clostridium perfringens has been studied in three strains shown to have high (NCTC8237), intermediate (strain 13) and low (NCTC8533) phospholipase C activity in the culture supernatant. The phospholipase C activity was shown to be related to cpa mRNA levels. Primer extension studies were performed to locate the cpa promoter regions in strains NCTC8237 and 13. Differences in promoter sequences could account for the differences in alpha-toxin production between strains 13 and NCTC8237. In contrast, the differences in alpha-toxin production between strains NCTC8237 and NCTC8533 were unlikely to be due to promoter differences because the upstream promoter-containing sequences were identical in these strains. The recombinant plasmid carrying the NCTC8237 cpa gene was introduced into strains 13 and NCTC8533. The level of production of the alpha-toxin was 16-fold higher in strain 13, indicating the presence of strain-dependant regulatory systems.

REGISTRY NUMBERS: 9001-86-9Q: PHOSPHOLIPASE C; 63551-76-8Q: PHOSPHOLIPASE C

DESCRIPTORS:

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics; Enzymology--Biochemistry and Molecular Biophysics; Genetics; Molecular Genetics--Biochemistry and Molecular Biophysics; Physiology; Toxicology

BIOSYSTEMATIC NAMES: Endospore-forming Gram-Positives--Eubacteria, Bacteria, Microorganisms

ORGANISMS: endospore-forming gram-positive rods and cocci (Endospore-forming Gram-Positives); Clostridium perfringens (Endospore-forming Gram-Positives)

COMMON TAXONOMIC TERMS: Bacteria; Eubacteria; Microorganisms

CHEMICALS & BIOCHEMICALS: PHOSPHOLIPASE C  
MISCELLANEOUS TERMS: ALPHA-TOXIN; GENE EXPRESSION; MOLECULAR GENETICS;  
PHOSPHOLIPASE C; PROMOTER MAPPING; STRAIN-NCTC8237; STRAIN-NCTC8533;  
STRAIN-13; TOXICOLOGY

CONCEPT CODES:

10064 Biochemistry studies - Proteins, peptides and amino acids  
10300 Replication, transcription, translation  
10808 Enzymes - Physiological studies  
22501 Toxicology - General and methods  
31000 Physiology and biochemistry of bacteria  
31500 Genetics of bacteria and viruses

BIOSYSTEMATIC CODES:

07810 Endospore-forming Gram-Positives

2/9/66 (Item 8 from file: 357)

DIALOG(R) File 357:Derwent Biotech Res.

(c) 2005 Thomson Derwent & ISI. All rts. reserv.

0080240 DBR Accession No.: 88-11089

**Isolation of promoter from 2 anaerobic bacteria - cloning Clostridium  
absonum and Bacteroides thetaiotaomicron DNA in Escherichia coli; DNA  
sequence**

AUTHOR: Roberts I; Hylemon P B; Holmes W M

CORPORATE SOURCE: Department of Microbiology, Medical College of Virginia,  
Virginia Commonwealth University, Richmond, Virginia 23298-0678, USA.

JOURNAL: Microbios (54, 219, 87-99) 1988

CODEN: MCBIA7

LANGUAGE: English

ABSTRACT: DNA fragments were cloned from 2 anaerobes, Clostridium absonum ATCC 27637 and Bacteroides thetaiotaomicron ATCC 29148 into Escherichia coli RR-1, using promoter probe plasmid pK01, carrying a beta-galactosidase (EC-3.2.1.23) gene (galK). About 10% of clones contained promoters functional in E. coli. Plasmid pRCL101, which directed expression of beta-galactosidase to 977 U, and had an insert of 88 bp, was sequenced. 5 Putative -10 sequences and 1 putative -35 sequence were found. The fragment originated from C. absonum DNA. Transcription in vitro showed that there were 2 transcription start sites near the end of the 88 bp sequence, and that C. absonum RNA-polymerase (EC-2.7.7.6) recognized the DNA fragment as an initiation point for transcription. A translation start site was found followed by an open reading frame. From a primer extension assay, 2 transcription initiation sites were found downstream from the 88 bp sequence. This method for promoter isolation may also assure efficient expression of genes in E. coli as well as Clostridium sp., a factor which may be important for large scale protein production in E.coli. (20 ref)

E.C. NUMBERS: 3.2.1.23; 2.7.7.6

DESCRIPTORS: anaerobe Clostridium absonum, Bacteroides thetaiotaomicron promoter, cloning, plasmid pRCL101, DNA sequence bacterium

SECTION: Microbiology-Genetics (A1)

2/9/67 (Item 9 from file: 357)

DIALOG(R) File 357:Derwent Biotech Res.

(c) 2005 Thomson Derwent & ISI. All rts. reserv.

0070964 DBR Accession No.: 88-01312

**Characterization of the tetanus toxin promoter and expression of nontoxic  
fragments in E. coli - RNA-polymerase isolation and characterization  
from Clostridium tetani and Escherichia coli (conference abstract)**

AUTHOR: Eisel U; Binz T; Niemann H

CORPORATE SOURCE: Institut fuer Medizinische Virologie, Frankfurter Str.  
107, D-6300 Giessen, Germany.

JOURNAL: Biol.Chem.Hoppe Seyler (368, 9, 1037-38) 1987

CODEN: BCHSEI

LANGUAGE: English

ABSTRACT: The tetanus toxin promoter has been characterized. RNA-polymerase (EC-2.7.7.6) was isolated from *Clostridium tetani* toxigenic strain E88, nontoxigenic variant EK11, and *Escherichia coli*. RNA-polymerase from EK11 was as efficient in transcription from the toxin promoter as that from strain E88. A region upstream from the transcription start point showed close homology with promoter sequences of other Gram-positive organisms and the *E. coli* lac promoter. The promoter region and various truncated forms were inserted upstream from the chloramphenicol-acetyltransferase (CAT) gene, and the CAT activity of *E. coli* lysates was studied to determine the strength of individual promoters. An A+T-rich region upstream of the toxin promoter was found to have an enhancing effect on the transcription rate in *E. coli*. Individual toxin-specific peptides spanning the whole toxin molecule have been expressed in *E. coli*. Fusion proteins with the MS-2 polymerase were synthesized using the pEx31 expression system, and the products were characterized with respect to immunogenicity. (1 ref)

E.C. NUMBERS: 2.7.7.6

DESCRIPTORS: *Clostridium tetani* tetanus toxin promoter characterization, RNA-polymerase isol., characterization, *Escherichia coli* bacterium enzyme EC-2.7.7.6

SECTION: Pharmaceuticals-Vaccines; Microbiology-Genetics (D4,A1)

2/9/68 (Item 1 from file: 51)

DIALOG(R) File 51:Food Sci.&Tech.Abs

(c) 2005 FSTA IFIS Publishing. All rts. reserv.

00758395 1998-05-c0510 SUBFILE: FSTA

**Identification and characterization of sporulation-dependent promoters upstream of the enterotoxin gene (cpe) of *Clostridium perfringens*.**

Yuling Zhao; Melville, S. B.

Correspondence (Reprint) address, S. B. Melville, Dep. of Microbiol. & Immunol., Univ. of Tennessee, Memphis, TN 38163, USA. Tel. (901) 448-6779. Fax (901) 448-8462. E-mail sbmelville(a)utmeml.utmem.edu

Journal of Bacteriology 1998 , 180 (1) 136-142

NOTE: 30 ref.

DOCUMENT TYPE: Journal Article ISSN: 0021-9193

LANGUAGE: English

In *Clostridium perfringens*, CPE enterotoxin synthesis is linked with sporulation. Although several apparent mRNA 5' ends have been identified in the region immediately upstream of the cpe gene, the promoters responsible for sporulation-dependent regulation of cpe have not been identified. To determine if these 5' ends represent actual promoter elements, a series of mutational and biochemical analyses of transcription of the upstream region were carried out. 3 promoter sites (P1-3) responsible for CPE synthesis were identified. DNA sequences upstream of P1 were similar to consensus SigK-dependent promoters, while P2 and P3 were similar to consensus SigE-dependent promoters. SigE and SigK are both sporulation-associated sigma factors which are active in the mother cell compartment of sporulating cells of *Bacillus subtilis*, the same compartment in which CPE is synthesized in *C. perfringens*.

DESCRIPTORS (HEADINGS): CLOSTRIDIUM; ENTEROTOXINS; FOOD SAFETY; GENETICS

DESCRIPTORS: GENES; PROMOTERS

SECTION HEADINGS: Hygiene & to


[ExPASy Home page](#)
[Site Map](#)
[Search ExPASy](#)
[Contact us](#)
[Proteomics tools](#)
[Swiss-Prot](#)

Search  for

=====

Welcome to the SIB BLAST Network Service

If results of this search are reported or published, please mention that the computation was performed at the SIB using the BLAST network service. The SIB BLAST network service uses a server developed at SIB and the NCBI BLAST 2 software.

In case of problems, please read the [online BLAST help](#).  
If your question is not covered, please contact [<helpdesk@expasy.org>](mailto:helpdesk@expasy.org).

NCBI BLAST program reference [PMID:9254694]:

Altschul S.F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W., Lipman D.J. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res. 25:3389-3402(1997).

=====

Query: 30 AA

Date run: 2005-02-15 12:40:58 UTC+0100 on sib-gml.unil.ch

Program: NCBI BLASTP 1.5.4-Paracel [2003-06-05]

Database: EXPASY/UniProt

1,794,555 sequences; 574,459,479 total letters

UniProt Release 4.1 consists of: Swiss-Prot Release 46.1 of 15-Feb-2005: 170140 en  
TrEMBL Release 29.1 of 15-Feb-2005: 1614107 entrie

[Taxonomic view](#)

[NiceBlast view](#)

[Printable view](#)

#### List of potentially matching sequences

Send selected sequences to

☐ Include query sequence


Db	AC	Description	Score	E-value
<input type="checkbox"/>	tr Q83U44	_CLOPE Beta2-toxin [cpb2] [Clostridium perfringens]	101	2e-21
<input type="checkbox"/>	tr O86264	_CLOPE Beta 2 toxin precursor [Clostridium perfringens C]	101	2e-21
<input type="checkbox"/>	tr Q93MD0	_CLOPE Beta2-toxin [cpb2] [Clostridium perfringens]	95	1e-19
<input type="checkbox"/>	tr Q8U1W2	_PYRFU Hypothetical protein PF1092 [PF1092] [Pyrococcus...]	32	1.1
<input type="checkbox"/>	sp P45963	ACR7_CAEEL Acetylcholine receptor, alpha-type subunit ...	32	1.5
<input type="checkbox"/>	tr Q81L65	_BACAN Iron compound ABC transporter, iron compound-bin...	32	1.5
<input type="checkbox"/>	tr Q731T5	_BACC1 HlyC domain protein [BCE4080] [Bacillus cereus (...]	32	1.5
<input type="checkbox"/>	tr Q8AWN0	_9SMEG Rhodopsin (Fragment) [Rhod] [Spinachia spinachia]	31	2.0

<input type="checkbox"/>	tr <a href="#">Q65ZZ8</a>	_BORG Hypothetical protein [BG0800] [Borrelia garinii]	<a href="#">31</a>	2.0
<input type="checkbox"/>	tr <a href="#">Q5NER8</a>	_FRATT Hypothetical protein [FTT1550] [Francisella tula...	<a href="#">31</a>	2.7
<input type="checkbox"/>	tr <a href="#">Q33865</a>	_BACPU Plasmid pSH1452, Rep [Bacillus pumilus (Bacillus...	<a href="#">30</a>	3.6
<input type="checkbox"/>	tr <a href="#">Q91F76</a>	_IRV6 450L [Chilo iridescent virus (CIV) (Insect irides...	<a href="#">30</a>	4.8
<input type="checkbox"/>	tr <a href="#">Q72MW1</a>	_LEPIC Hypothetical protein [LIC13078] [Leptospira inte...	<a href="#">30</a>	4.8
<input type="checkbox"/>	tr <a href="#">Q6LF23</a>	_PLAF7 Hypothetical protein [PFF1215w] [Plasmodium falc...	<a href="#">30</a>	4.8
<input type="checkbox"/>	tr <a href="#">Q73K80</a>	_TREDE FMN-binding domain protein [TDE2340] [Treponema ...	<a href="#">29</a>	6.4
<input type="checkbox"/>	tr <a href="#">Q8AWP8</a>	_9TELE Rhodopsin (Fragment) [Rhod] [Cerattias holboelli]	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q8AWL3</a>	_PHOGU Rhodopsin (Fragment) [Rhod] [Pholis gunnellus (B...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q8AWJ5</a>	_9PERO Rhodopsin (Fragment) [Rhod] [Mene maculata]	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q69BZ0</a>	_CYCLU Rhodopsin (Fragment) [Rhod] [Cyclopterus lumpus ...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q69BX5</a>	_9PERC Rhodopsin (Fragment) [Rhod] [Triacanthodes sp. A...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q46102</a>	_CAMJE CdtC (Cytolethal distending toxin C) [cdtC] [Cam...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q64TC5</a>	_BACFR Hypothetical protein [BF2505] [Bacteroides fragi...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q8A943</a>	_BACTN Hypothetical protein [BT0974] [Bacteroides theta...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q5HX88</a>	_CAMJE Cytolethal distending toxin, subunit C [cdtC] [C...	<a href="#">29</a>	8.6
<input type="checkbox"/>	tr <a href="#">Q7B3S5</a>	_CAMJE CdtC protein (Fragment) [cdtC] [Campylobacter je...	<a href="#">29</a>	8.6

## Graphical overview of the alignments

 Click here

to resubmit your query after masking regions matching [PROSITE](#) profiles or [Pfam](#) HMMs


( [Help](#)) (use [ScanProsite](#) for more details about PROSITE matches)

## Profile hits

=====

## Pfam hits

=====



## Alignments

tr Q83U44 Beta2-toxin [cpb2] [Clostridium perfringens] 265 AA  
 Q83U44\_CLOPE [align](#)

Score = 101 bits (231), Expect = 2e-21  
 Identities = 30/30 (100%), Positives = 30/30 (100%)

Query: 1 MKKIISKFTVIFMFSCFLIVGAISPMKASA 30  
 MKKIISKFTVIFMFSCFLIVGAISPMKASA  
 Sbjct: 1 MKKIISKFTVIFMFSCFLIVGAISPMKASA 30

tr Q86264 Beta 2 toxin precursor [Clostridium perfringens C] 265 AA  
 Q86264\_CLOPE [align](#)

Score = 101 bits (231), Expect = 2e-21  
 Identities = 30/30 (100%), Positives = 30/30 (100%)

Query: 1 MKKIISKFTVIFMFSCFLIVGAISPMKASA 30  
 MKKIISKFTVIFMFSCFLIVGAISPMKASA  
 Sbjct: 1 MKKIISKFTVIFMFSCFLIVGAISPMKASA 30

tr Q93MD0 **Beta2-toxin [cpb2] [Clostridium perfringens]** 265 AA  
Q93MD0\_CLOPE align

Score = 95.2 bits (217), Expect = 1e-19  
Identities = 29/30 (96%), Positives = 29/30 (96%)

Query: 1 MKKIISKFTVIFMFSCFLIVGAISPMKASA 30  
MKKIISKFTVIFMFS FLIVGAISPMKASA  
Sbjct: 1 MKKIISKFTVIFMFSYFLIVGAISPMKASA 30

tr Q8U1W2 **Hypothetical protein PF1092 [PF1092] [Pyrococcus** 389  
Q8U1W2\_PYRFU **furiosus]** AA  
align

Score = 32.0 bits (68), Expect = 1.1  
Identities = 9/12 (75%), Positives = 9/12 (75%)

Query: 4 IISKFTVIFMFS 15  
II F VIFMFS  
Sbjct: 99 IIATFPVIFMFS 110

sp P45963 **Acetylcholine receptor, alpha-type subunit acr-7 precursor** 538  
ACR7\_CAEEL **[acr-7]** AA  
**[Caenorhabditis elegans]** align

Score = 31.6 bits (67), Expect = 1.5  
Identities = 12/24 (50%), Positives = 14/24 (58%), Gaps = 3/24 (12%)

Query: 4 IISKFTVIFMFSCFLIVGAISPMK 27  
II FT++F CF V AI P K  
Sbjct: 515 II--FTIVFIICCFIFV-AIPPIK 535

tr Q81L65 **Iron compound ABC transporter, iron compound-binding** 324  
Q81L65\_BACAN **protein** AA  
**[BA4766] [Bacillus anthracis]** align

Score = 31.6 bits (67), Expect = 1.5  
Identities = 9/14 (64%), Positives = 11/14 (78%)

Query: 1 MKKIISKFTVIFMF 14  
MKKI S F V+F+F  
Sbjct: 1 MKKILSIFIVVFLF 14

tr Q731T5 **HlyC domain protein [BCE4080] [Bacillus cereus (strain** 99 AA  
Q731T5\_BACC1 **ATCC**  
**10987)]** align



Score = 31.6 bits (67), Expect = 1.5  
Identities = 14/26 (53%), Positives = 14/26 (53%), Gaps = 9/26 (34%)

Query: 2 KKIISK-----FTVI-FMFSCFLIV 20  
KKI K F VI FMFSC IV  
Sbjct: 60 KKIFPKKYIIEIFRVIVFMFSC--IV 83

tr Q8AWN0 Rhodopsin (Fragment) [Rhod] [Spinachia spinachia] 253 AA  
Q8AWN0\_9SMEG

[align](#)

Score = 31.2 bits (66), Expect = 2.0  
Identities = 10/18 (55%), Positives = 11/18 (60%), Gaps = 6/18 (33%)

Query: 10 VIFMFSC-FLIVGAISPM 26  
VI+MF C FLI PM  
Sbjct: 160 VIYMFTCHFLI-----PM 172

tr Q65ZZ8 Hypothetical protein [BG0800] [Borrelia garinii] 186 AA  
Q65ZZ8\_BORGA

[align](#)

Score = 31.2 bits (66), Expect = 2.0  
Identities = 10/17 (58%), Positives = 11/17 (63%), Gaps = 3/17 (17%)

Query: 1 MKKIISKFTVIFMFSCF 17  
M KI SKF F+F CF  
Sbjct: 1 MNKILSKF---FLFFCF 14

tr Q5NER8 Hypothetical protein [FTT1550] [Francisella tularensis] 180  
Q5NER8\_FRATT (subsp. tularensis) AA  
[align](#)

Score = 30.8 bits (65), Expect = 2.7  
Identities = 11/20 (55%), Positives = 11/20 (55%), Gaps = 4/20 (20%)

Query: 1 MKKIISKFTVIFM----FSC 16  
MKK ISK VI M F C  
Sbjct: 1 MKKLISKIGVIIMALGLFGC 20

tr O33865 Plasmid pSH1452, Rep [Bacillus pumilus (Bacillus] 177  
O33865\_BACPU mesentericus) AA  
[align](#)

Score = 30.3 bits (64), Expect = 3.6  
Identities = 10/20 (50%), Positives = 11/20 (55%), Gaps = 7/20 (35%)

Query: 2 KKIIS-----KFTVIFMF 14  
KKIIS F V+FMF  
Sbjct: 4 KKIISLITILVLTFSVVFMF 23

tr Q91F76 450L [Chilo iridescent virus (CIV) (Insect iridescent virus type 6)] 74 AA  
align

Score = 29.9 bits (63), Expect = 4.8  
Identities = 9/14 (64%), Positives = 9/14 (64%), Gaps = 4/14 (28%)

Query: 1 MKKIISKFTVIFMF 14  
MKKI T IFMF  
Sbjct: 1 MKKI---TMIFMF 10

tr Q72MW1 Hypothetical protein [LIC13078] [Leptospira interrogans 289 AA  
Q72MW1\_LEPIC (serogroup Icterohaemorrhagiae / serovar Copenhageni)]  
align

Score = 29.9 bits (63), Expect = 4.8  
Identities = 10/12 (83%), Positives = 10/12 (83%)

Query: 1 MKKIISKFTVIF 12  
MKKIIS F VIF  
Sbjct: 15 MKKIISLFFVIF 26

tr Q6LF23 Hypothetical protein [PFF1215w] [Plasmodium falciparum 413  
Q6LF23\_PLAF7 (isolate 3D7)] AA  
align

Score = 29.9 bits (63), Expect = 4.8  
Identities = 12/22 (54%), Positives = 14/22 (63%), Gaps = 6/22 (27%)

Query: 4 IISKFTVIFMFSCF----LIVG 21  
II KF +IF FSCF I+G  
Sbjct: 243 II-KF-IIFTFSCFIYAIIIIIG 262

tr Q73K80 FMN-binding domain protein [TDE2340] [Treponema 120  
Q73K80\_TREDE denticola] AA  
align

Score = 29.5 bits (62), Expect = 6.4  
Identities = 11/22 (50%), Positives = 13/22 (59%), Gaps = 8/22 (36%)

Query: 1 MKKIISKFTVI-----FMFSC 16  
MKKI FT+I F+FSC  
Sbjct: 1 MKKIC--FTIIVFALSIFLFSC 20

tr Q8AWP8 Rhodopsin (Fragment) [Rhod] [Ceratias holboelli] 253 AA  
Q8AWP8\_9TELE

align

Score = 29.1 bits (61), Expect = 8.6

Identities = 8/11 (72%), Positives = 10/11 (90%), Gaps = 1/11 (9%)

Query: 10 VIFMFSC-FLI 19  
VI+MFSC FL+  
Sbjct: 160 VIYMFSCFLV 170

tr Q8AWL3 Rhodopsin (Fragment) [Rhod] [Pholis gunnellus] 249  
Q8AWL3\_PHOGU (Butterfish) (Rock  
gunnel)] AA  
align

Score = 29.1 bits (61), Expect = 8.6

Identities = 8/11 (72%), Positives = 9/11 (81%), Gaps = 1/11 (9%)

Query: 10 VIFMFSC-FLI 19  
VI+MF C FLI  
Sbjct: 156 VIYMFCTHFLI 166

tr Q8AWJ5 Rhodopsin (Fragment) [Rhod] [Mene maculata] 253 AA  
Q8AWJ5\_9PERO align

Score = 29.1 bits (61), Expect = 8.6

Identities = 10/16 (62%), Positives = 11/16 (68%), Gaps = 5/16 (31%)

Query: 10 VIFMFSC-FL----IV 20  
VI+MFSC FL IV  
Sbjct: 160 VIYMFSCFLTPLTIV 175

tr Q69BZ0 Rhodopsin (Fragment) [Rhod] [Cyclopterus lumpus] 252  
Q69BZ0\_CYCLU (Lumpsucker)] AA  
align

Score = 29.1 bits (61), Expect = 8.6

Identities = 8/11 (72%), Positives = 9/11 (81%), Gaps = 1/11 (9%)

Query: 10 VIFMFSC-FLI 19  
VI+MF C FLI  
Sbjct: 160 VIYMFCTHFLI 170

tr Q69BX5 Rhodopsin (Fragment) [Rhod] [Triacanthodes sp. AD-2003] 216 AA

Q69BX5\_9PERC

[align](#)

Score = 29.1 bits (61), Expect = 8.6

Identities = 10/16 (62%), Positives = 11/16 (68%), Gaps = 5/16 (31%)

Query: 10 VIFMFSC-FL----IV 20

VI+MFSC FL IV

Sbjct: 123 VIYMFSCFLTPLTIV 138

tr Q46102 CdtC (Cytolethal distending toxin C) [cdtC]  
Q46102\_CAMJE [Campylobacter  
jejuni]

189

AA

[align](#)

Score = 29.1 bits (61), Expect = 8.6

Identities = 9/14 (64%), Positives = 9/14 (64%), Gaps = 3/14 (21%)

Query: 1 MKKIISKFTVIFMF 14

MKKII T FMF

Sbjct: 1 MKKII---TLFFMF 11

tr Q64TC5 Hypothetical protein [BF2505] [Bacteroides fragilis] 161 AA  
Q64TC5\_BACFR

[align](#)

Score = 29.1 bits (61), Expect = 8.6

Identities = 11/18 (61%), Positives = 11/18 (61%), Gaps = 4/18 (22%)

Query: 9 TVIFMFSCFLIVGAISPM 26

TVIFM L VGA PM

Sbjct: 135 TVIFM---LAVGATFPM 148

tr Q8A943 Hypothetical protein [BT0974] [Bacteroides  
Q8A943\_BACTN thetaiotaomicron]

164

AA

[align](#)

Score = 29.1 bits (61), Expect = 8.6

Identities = 11/18 (61%), Positives = 11/18 (61%), Gaps = 4/18 (22%)

Query: 9 TVIFMFSCFLIVGAISPM 26

TVIFM L VGA PM

Sbjct: 138 TVIFM---LAVGATFPM 151

tr Q5HX88 Cytolethal distending toxin, subunit C [cdtC]  
Q5HX88\_CAMJE [Campylobacter  
jejuni RM1221]

189

AA

[align](#)

Score = 29.1 bits (61), Expect = 8.6

Identities = 9/14 (64%), Positives = 9/14 (64%), Gaps = 3/14 (21%)

Query: 1 MKKIISKFTVIFMF 14  
MKKII T FMF  
Sbjct: 1 MKKII---TLFFMF 11

tr Q7B3S5 CdtC protein (Fragment) [cdtC] [Campylobacter jejuni] 111 AA  
Q7B3S5\_CAMJE  
align

Score = 29.1 bits (61), Expect = 8.6  
Identities = 9/14 (64%), Positives = 9/14 (64%), Gaps = 3/14 (21%)

Query: 1 MKKIISKFTVIFMF 14  
MKKII T FMF  
Sbjct: 1 MKKII---TLFFMF 11

Database: EXPASY/UniProt  
Posted date: Feb 15, 2005 11:36 AM  
Number of letters in database: 574,459,479  
Number of sequences in database: 1,794,555

Lambda K H  
0.343 0.274 1.80

Gapped  
Lambda K H  
0.294 0.110 0.610

Matrix: PAM30  
Gap Penalties: Existence: 9, Extension: 1  
length of query: 30  
length of database: 574,459,479  
effective HSP length: 21  
effective length of query: 9  
effective length of database: 536,773,824  
effective search space: 4830964416  
effective search space used: 4830964416  
T: 16  
A: 40  
X1: 15 ( 7.4 bits)  
X2: 35 (14.8 bits)  
X3: 58 (24.6 bits)  
S1: 40 (21.6 bits)  
S2: 61 (29.1 bits)

Wallclock time: 2 seconds

 <a href="#">ExPASy Home page</a>	<a href="#">Site Map</a>	<a href="#">Search ExPASy</a>	<a href="#">Contact us</a>	<a href="#">Proteomics tools</a>	<a href="#">Swiss-Prot</a>
--	--------------------------	-------------------------------	----------------------------	----------------------------------	----------------------------

[ExPASy Home page](#)[Site Map](#)[Search ExPASy](#)[Contact us](#)[Swiss-Prot](#)Search  for   

# NiceProt

## View of

## TrEMBL:

## O86264

[Printer-friendly view](#)[Request update](#)[Quick BlastP search](#)

[\[Entry info\]](#) [\[Name and origin\]](#) [\[References\]](#) [\[Comments\]](#) [\[Cross-references\]](#) [\[Keywords\]](#)  
[\[Features\]](#) [\[Sequence\]](#) [\[Tools\]](#)

*Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.*

### Entry information

Entry name **O86264\_CLOPE**  
 Primary accession number **O86264**  
 Secondary accession numbers None  
 Entered in TrEMBL in Release 08, November 1998  
 Sequence was last modified in Release 08, November 1998  
 Annotations were last modified in Release 24, June 2003

### Name and origin of the protein

Protein name **Beta 2 toxin [Precursor]**  
 Synonyms None  
 Gene name None  
 From *Clostridium perfringens* C [TaxID: 79668]  
 Taxonomy Bacteria; Firmicutes; Clostridia; Clostridiales; Clostridiaceae; Clostridium.

### References

#### [1] NUCLEOTIDE SEQUENCE.

**STRAIN=CWC245;**  
 DOI=10.1016/S0378-1119(97)00493-9;MEDLINE=98085977;PubMed=9426008 [NCBI, ExPASy, EBI, Israel, Japan]  
 Gibert M., Jolivet-Reynaud C., Popoff M.R.;  
 "Beta2 toxin, a novel toxin produced by *Clostridium perfringens*."; *Gene* 203:65-73(1997).

#### [2] NUCLEOTIDE SEQUENCE.

**STRAIN=CWC245;**  
 Popoff M.R.;  
 Submitted (JAN-1998) to the EMBL/GenBank/DDBJ databases.

### Comments

None


### Cross-references

EMBL L77965; AAC27654.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence]  
 PIR JC6515; JC6515.

ProDom [Domain structure / List of seq. sharing at least 1 domain]  
 HOBACGEN [Family / Alignment / Tree]  
 ProtoMap O86264.  
 PRESAGE O86264.  
 ModBase O86264.  
 SMR O86264; 8972E69F52B26CBD.  
 SWISS-2DPAGE Get region on 2D PAGE.  
 UniRef View cluster of proteins with at least 50% / 90% identity.

**Keywords**

Signal.

**Features**

Feature table viewer

Key	From	To	Length	Description
SIGNAL	1	30	30	Potential.
CHAIN	31	265	235	beta 2 toxin.

**Sequence information**

Length: **265 AA** [This is the length of the unprocessed precursor]  
 Molecular weight: **30963 Da** [This is the MW of the unprocessed precursor]

CRC64: **8972E69F52B26CBD** [This is a checksum on the sequence]

<u>10</u>	<u>20</u>	<u>30</u>	<u>40</u>	<u>50</u>	<u>60</u>
MKKIISKFTV	IFMFSCFLIV	GAISPMKASA	KEIDAYRKVM	ENYLNALKNY	DINTVVNISE
<u>70</u>	<u>80</u>	<u>90</u>	<u>100</u>	<u>110</u>	<u>120</u>
DERVNNVEQY	REMLEDFKYD	PNQQLKSFEI	LNSQKSDNKE	IFNVKTEFLN	GAIYDMEFTV
<u>130</u>	<u>140</u>	<u>150</u>	<u>160</u>	<u>170</u>	<u>180</u>
SSKDGKLIVS	DMERTKVENE	GKYILTPSFR	TQVCTWDEL	AQAIGGVYPQ	TYSRFTYYA
<u>190</u>	<u>200</u>	<u>210</u>	<u>220</u>	<u>230</u>	<u>240</u>
DNILLNFRQY	ATGSRDLKV	EYSVVDHWMW	KDDVKASQMV	YGQNPDSARQ	IRLYIEKGQS
<u>250</u>	<u>260</u>				
FYKYRIRIKN	FTPASIRVFG	EGYCA			

O86264 in FASTA format

*View entry in original TrEMBL format*

*View entry in raw text format (no links)*

*Request for annotation of this TrEMBL entry*

BLAST submission on  
 ExPASy/SIB  
 or at NCBI (USA)




Sequence analysis tools: ProtParam, ProtScale,  
 Compute pI/Mw, PeptideMass, PeptideCutter,  
 Dotlet (Java)



ScanProsite, MotifScan



Search the SWISS-MODEL Repository

 <b>ExPASy Home page</b>		<b>Site Map</b>		<b>Search ExPASy</b>		<b>Contact us</b>		<b>Swiss-Prot</b>	
Hosted by NCSC US	Mirror sites:	Australia	Bolivia	Brazil <small>new</small>	Canada	China	Korea	Switzerland	Taiwan



SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1951-2005/Feb W2

(c) format only 2005 The Dialog Corp.

\*File 155: Medline has resumed updating. Please see  
HELP NEWS 155 for details.

File 5:Biosis Previews(R) 1969-2005/Feb W1

(c) 2005 BIOSIS

\*File 5: Price change effective Jan 1, 2005. Enter HELP  
RATES 5 for details.

File 390:Beilstein Facts Nov. 2004

(c) 2005 Beilstein GmbH

\*File 390: File has been reloaded. Please see HELP NEWS 390.  
IMPORTANT - Price based on output. See HELP RATES 390.

File 399:CA SEARCH(R) 1967-2005/UD=14207

(c) 2005 American Chemical Society

\*File 399: Use is subject to the terms of your user/customer agreement.  
Alert feature enhanced for multiple files, etc. See HELP ALERT.

File 34:SciSearch(R) Cited Ref Sci 1990-2005/Feb W1

(c) 2005 Inst for Sci Info

\*File 34: Price change effective Jan 1, 2005. Enter HELP  
RATES 34 for details.

File 73:EMBASE 1974-2005/Feb W1

(c) 2005 Elsevier Science B.V.

\*File 73: Price change effective Jan 1, 2005. Enter HELP  
RATES 73 for details.

File 144:Pascal 1973-2005/Feb W1

(c) 2005 INIST/CNRS

\*File 144: Price change effective Jan 1, 2005. Enter HELP  
RATES 144 for details.

File 357:Derwent Biotech Res. 1982-2005/Feb W2

(c) 2005 Thomson Derwent & ISI

File 71:ELSEVIER BIOBASE 1994-2005/Feb W1

(c) 2005 Elsevier Science B.V.

File 143:Biol. & Agric. Index 1983-2005/Jan

(c) 2005 The HW Wilson Co

File 51:Food Sci.&Tech.Abs 1969-2005/Feb W2

(c) 2005 FSTA IFIS Publishing

File 156:ToxFile 1965-2005/Feb W1

(c) format only 2005 The Dialog Corporation

\*File 156: Updating of ToxFile has resumed, with  
UD=20041205.

File 484:Periodical Abs Plustext 1986-2005/Feb W1

(c) 2005 ProQuest

\*File 484: SELECT IMAGE AVAILABILITY FOR PROQUEST FILES  
ENTER 'HELP PROQUEST' FOR MORE

File 98:General Sci Abs/Full-Text 1984-2004/Sep

(c) 2004 The HW Wilson Co.

File 434:SciSearch(R) Cited Ref Sci 1974-1989/Dec

(c) 1998 Inst for Sci Info

\*File 434: Price change effective Jan 1, 2005. Enter HELP  
RATES 434 for details.

File 453:Drugs of the Future 1990-2002/Oct

(c) 2002 Prous Science

\*File 453: Updating of this file has temporarily ceased due to  
a production system change.

File 286:Biotechnology Directory Current Jan B2

(c) 2005 BioCommerce Data Ltd.

\*File 286: Price change effective Jan 1, 2005.  
Enter HELP RATES286 after 1/1/2005 for more information.

File 8:Ei Compendex(R) 1970-2005/Jan W3

(c) 2005 Elsevier Eng. Info. Inc.

\*File 8: Price change effective Jan 1, 2005. Enter HELP  
RATES 8 for details.

File 35:Dissertation Abs Online 1861-2005/Jan

(c) 2005 ProQuest Info&Learning

File 50:CAB Abstracts 1972-2005/Jan

(c) 2005 CAB International

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1951-2005/Feb W2

(c) format only 2005 The Dialog Corp.

\*File 155: Medline will be reloaded shortly and accession numbers will change.

File 5:Biosis Previews(R) 1969-2005/Feb W2

(c) 2005 BIOSIS

\*File 5: Price change effective Jan 1, 2005. Enter HELP RATES 5 for details.

File 34:SciSearch(R) Cited Ref Sci 1990-2005/Feb W2

(c) 2005 Inst for Sci Info

\*File 34: Price change effective Jan 1, 2005. Enter HELP RATES 34 for details.

File 35:Dissertation Abs Online 1861-2005/Jan

(c) 2005 ProQuest Info&Learning

File 48:SPORTDiscus 1962-2005/May

(c) 2005 Sport Information Resource Centre

File 65:Inside Conferences 1993-2005/Feb W2

(c) 2005 BLDSC all rts. reserv.

File 71:ELSEVIER BIOBASE 1994-2005/Feb W1

(c) 2005 Elsevier Science B.V.

File 73:EMBASE 1974-2005/Feb W1

(c) 2005 Elsevier Science B.V.

\*File 73: Price change effective Jan 1, 2005. Enter HELP RATES 73 for details.

File 91:MANTIS(TM) 1880-2005/Feb

2001 (c) Action Potential

File 94:JICST-EPlus 1985-2005/Jan W1

(c)2005 Japan Science and Tech Corp(JST)

File 98:General Sci Abs/Full-Text 1984-2004/Sep

(c) 2004 The HW Wilson Co.

File 135:NewsRx Weekly Reports 1995-2005/Feb W1

(c) 2005 NewsRx

\*File 135: New newsletters are now added. See Help News135 for the complete list of newsletters.

File 144:Pascal 1973-2005/Feb W1

(c) 2005 INIST/CNRS

\*File 144: Price change effective Jan 1, 2005. Enter HELP RATES 144 for details.

File 149:TGG Health&Wellness DB(SM) 1976-2005/Feb W1

(c) 2005 The Gale Group

File 156:ToxFile 1965-2005/Feb W2

(c) format only 2005 The Dialog Corporation

\*File 156: Updating of ToxFile has resumed, with UD=20041205.

File 159:Cancerlit 1975-2002/Oct

(c) format only 2002 Dialog Corporation

\*File 159: Cancerlit is no longer updating.

Please see HELP NEWS159.

File 162:Global Health 1983-2005/Jan

(c) 2005 CAB International

File 164:Allied & Complementary Medicine 1984-2005/Feb

(c) 2005 BLHCIS

File 172:EMBASE Alert 2005/Feb W1

(c) 2005 Elsevier Science B.V.

\*File 172: Price change effective Jan 1, 2005. Enter HELP RATES 172 for details.

File 266:FEDRIP 2004/Nov

Comp & dist by NTIS, Intl Copyright All Rights Res

File 369:New Scientist 1994-2005/Jan W5

(c) 2005 Reed Business Information Ltd.

File 370:Science 1996-1999/Jul W3

(c) 1999 AAAS

\*File 370: This file is closed (no updates). Use File 47 for more current information.

File 399:CA SEARCH(R) 1967-2005/UD=14207

(c) 2005 American Chemical Society

210

215

220

Pro Asp Ser Ala Arg Gln Ile Arg Leu Tyr Ile Glu Lys Gly Gln Ser  
225 230 235 240

Phe Tyr Lys Tyr Arg Ile Arg Ile Lys Asn Phe Thr Pro Ala Ser Ile  
245 250 255

Arg Val Phe Gly Glu Gly Tyr Cys Ala  
260 265

<210> 3  
<211> 267  
<212> DNA  
<213> Clostridium perfringens

267AA

<400> 3  
atttgggata tcttaaattt agcacagaag aatgttttaa tgaaataaag ataataaaaa 60  
gatatattaa ttatatagct gaaaatttat aattatatga taagtatagt taataaataa 120  
aaagtgttct cgggggacac ttttttgttt taaaaaggaa aatataaata aaatttagat 180  
aaaagtgtaa aataattatt tttatttttaa atttggttaa aatttgatat aattgaattg 240  
taaaaaaaat ttcagggggg aatataa 267

<210> 4  
<211> 90  
<212> DNA  
<213> Clostridium perfringens

30aa

90NA

<220>  
<221> CDS  
<222> (1)..(90)

<400> 4  
atg aaa aaa att att tca aag ttt act gta att ttt atg ttt tca tgt 48  
Met Lys Lys Ile Ile Ser Lys Phe Thr Val Ile Phe Met Phe Ser Cys  
1 5 10 15  
ttt ctt att gtt gga gca ata agt cca atg aaa gca agt gca 90  
Phe Leu Ile Val Gly Ala Ile Ser Pro Met Lys Ala Ser Ala  
20 25 30

<210> 5  
<211> 30  
<212> PRT  
<213> Clostridium perfringens

<400> 5

Met Lys Lys Ile Ile Ser Lys Phe Thr Val Ile Phe Met Phe Ser Cys  
1 5 10 15

tr Q46102 **CdtC (Cytolethal distending toxin C) [cdtC]**  
Q46102\_CAMJE **[Campylobacter**  
**jejuni]**

189  
AA  
align

Score = 29.1 bits (61), Expect = 8.6

Identities = 9/14 (64%), Positives = 9/14 (64%), Gaps = 3/14 (21%)

Query: 1 MKKIISKFTVIFMF 14  
MKKII T FMF  
Sbjct: 1 MKKII---TLFFMF 11